CEREBRAL PALSY
IS THE MOST
COMMON
PHYSICAL
DISABILITY OF
CHILDHOOD





ONLY 25% OF
INFANTS WITH
CEREBRAL PALSY
ARE DIAGNOSED
UNDER 6
MONTHS OFAGE





THE AVERAGE AGE OF DIAGNOSIS FOR CEREBRAL PALSY IN AUSTRALIA IS BETWEEN 12 AND 24 MONTHS OF AGE **GUIDELINES** 



MOST PARENTS
SUSPECT THE
DIAGNOSIS OF
CEREBRAL PALSY
BEFORE IT IS
FORMALLY GIVEN





A DELAY IN THE DIAGNOSIS OF CEREBRAL PALSY RESULTS IN DELAYED ACCESS TO INTERVENTION DURING AN IMPORTANT TIME OF BRAIN DEVELOPMENT 12

**GUIDELINES** 



NEW CLINICAL GUIDELINES ARE CHANGING THE STANDARD OF CARE FOR CHILDRENWITH CEREBRAL PALSY AND THEIR FAMILIES ACROSS AUSTRALIA

**GUIDELINES** 



NEW GUIDELINES CONTAIN 12 RECOMMENDATIONS FOR PRACTITIONERS THAT SUPPORTTHE EARLY DETECTION AND DIAGNOSIS OF CEREBRAL PALSY

**GUIDELINES** 

